

Quenched Light Hadron Spectrum and Decay Constants using Improved Wilson Fermion Actions

UKQCD Collaboration, presented by Richard Kenway^{a*}

^aDepartment of Physics & Astronomy, The University of Edinburgh, Edinburgh EH9 3JZ, Scotland

We compare results obtained using the Sheikholeslami-Wohlert (SW) fermion action with tree-level and tadpole-improved coefficients for $5.7 \leq \beta \leq 6.2$.

1. LATTICE PARAMETERS

The use of a tadpole-improved coefficient [1] for the clover term in the SW fermion action,

$$S_F = \sum_{xy} \bar{q}(x) \left\{ \left[1 - \frac{ic\kappa}{2} \sigma_{\mu\nu} F_{\mu\nu}(x) \right] \delta_{xy} - \kappa \left[(1 - \gamma_\mu) U_\mu(x) \delta_{x+\hat{\mu},y} + (1 + \gamma_\mu) U_\mu^\dagger(y) \delta_{x-\hat{\mu},y} \right] \right\} q(y), \quad (1)$$

where $c = 1/u_0^3$, $u_0 = \langle \frac{1}{3} \text{Tr} \square \rangle^{1/4}$, can be regarded as a step towards full $O(a)$ improvement [2]. We compare the resulting light hadron quantities with those obtained using the tree-level coefficient, $c = 1$, for which discretisation errors are $O(g^2 a)$ in perturbation theory [3], and also with data from GF11 [4] corresponding to $c = 0$, at a range of β values, to seek indications of better scaling behaviour.

Our data set comprises: at $\beta = 5.7$, Jacobi-smearred quark propagators at two κ values with $c = 1/u_0^3$ and $c = 1$ on $142 \times 16^3 \times 32$ configurations; at $\beta = 6.0$, fuzzed propagators [5] at three κ values with $c = 1/u_0^3$ on $499 \times 16^3 \times 48$ configurations; and, at $\beta = 6.2$, fuzzed (local) propagators at three (two) κ values with $c = 1/u_0^3$ ($c = 1$) on 130 (60) $24^3 \times 48$ configurations. We construct meson correlators from all κ combinations, but baryon correlators only from degenerate combinations. Consequently, we do not have enough baryon data to perform a reliable chiral extrapolation. Hadron masses are obtained from multi-exponential fits to various combinations of

smearred and local correlators. The results presented are from a preliminary analysis of our current data (see also [6]); higher statistics at $\beta = 6.2$ will be available soon.

2. HADRON SPECTRUM

The ratio of the nucleon to vector meson mass, m_N/m_V , at a fixed ratio of the pseudoscalar to vector meson mass, m_{PS}/m_V , has a noticeable dependence on c at our largest lattice spacing. The trend is towards improved scaling for this ratio as c is increased to the tadpole-improved value. Evidence for the latter is given in the Edinburgh plot in Figure 1.

Linear extrapolation of data for the pseudoscalar meson mass to $m_{PS}(\kappa_c, \kappa_c) = 0$ for the tadpole-improved action gives $u_0 \kappa_c = 0.12347(3)$, $0.12224(1)$, and $0.12208(2)$, for $\beta = 5.7$, 6.0 and 6.2 respectively, in reasonable agreement with the tree-level value of 0.125 . We define κ_{ud} , corresponding to degenerate u and d quarks, at $m_{PS}/m_V = m_\pi/m_\rho$. Then, as shown in Figure 2, our estimates for m_ρ in units of the square root of the string tension, $\sqrt{K}$, have decreasing dependence on the lattice spacing, a , as c is increased from 0 to $1/u_0^3$, although the dependence is not removed entirely.

The κ value corresponding to the strange quark mass, κ_s , may be fixed from any one of the ratios m_K/m_ρ , m_ϕ/m_ρ and m_{K^*}/m_ρ . In Figure 3 we show the results of fixing one of the first two ratios and calculating the third. Evidently, the K^* mass scales with the ρ mass in both cases, independently of whether $c = 1$ or $1/u_0^3$, but dif-

*Work supported by EPSRC grant GR/K41663 and PPARC grants GR/J98202, GR/K54601, GR/K55745.




Figure 1. Scaling at fixed m_{PS}/m_V for $c = 1/u_0^3$.

ferent values are obtained for m_{K^*}/m_ρ . This is an indication that the strange quark mass cannot be determined consistently in the quenched approximation [7]. We also find that J [8] is not changed by tadpole improvement, and scales at a value inconsistent with experiment.

Finally, we observe that, at all three β values, the magnitude of the meson spin splitting, $m_V^2 - m_{PS}^2$ in units of m_K or m_ρ , is insensitive to whether $c = 1$ or $1/u_0^3$.

3. DECAY CONSTANTS

Our quark propagators are not ‘rotated’, so $O(a)$ -improved matrix elements are constructed from improved quark fields:

$$\begin{aligned} q^I &= \sqrt{2\kappa u_0} \left[1 - \frac{\vec{D}}{2u_0} \right] q \\ &= \sqrt{2\kappa u_0} \left[1 + \frac{1}{2u_0} \left(\frac{1}{2\kappa} - \frac{1}{2\kappa_c} \right) \right] q \end{aligned} \quad (2)$$

where we have used the equations of motion and employed the tadpole-improvement prescription [1]. Thus, we take the tadpole-improved pion decay matrix element to be [9]

$$\langle 0 | \bar{q}^I \gamma_4 \gamma_5 q^I | \pi \rangle = Z_A 2\kappa u_0 \left[1 + \frac{1}{u_0} \left(\frac{1}{2\kappa} - \frac{1}{2\kappa_c} \right) \right]$$




Figure 2. m_ρ in units of the string tension.

$$\times \langle 0 | \bar{q} \gamma_4 \gamma_5 q | \pi \rangle \quad (3)$$

where we obtain κ_c from $m_{PS}(\kappa_c, \kappa_c) = 0$ and the current normalisation, Z_A , from tadpole-improved one-loop perturbation theory.

Comparison of the pion decay constant values obtained with different values of c is complicated by systematic effects inherent in the current normalisation prescription. Given this caveat, we show our tadpole-improved results along with those of GF11 [4] in Figure 4. Both sets of data suggest scaling may set in above $\beta = 6.0$.

The ratio of decay constants, f_K/f_π , is independent of the current normalisation and so may be determined more reliably. Our results, shown in Figure 5, are insensitive to whether $c = 1$ or $1/u_0^3$. For $\beta \geq 6.0$ the ratio agrees with experiment, but a weak lattice spacing dependence remains below this β value.

4. CONCLUSIONS

Using a tadpole-improved SW fermion action in quenched QCD, we conclude the following.

1. For $\beta \geq 5.7$, $m_\rho/\sqrt{K}$ has a weaker dependence on the lattice spacing (Figure 2), m_N/m_V and m_{K^*}/m_ρ scale (Figures 1 and 3), although




Figure 3. m_{K^*}/m_ρ for two definitions of κ_s .




Figure 4. f_π in units of m_ρ .

the latter clearly shows that the strange quark mass cannot be determined consistently.

2. Some quantities, such as m_{K^*}/m_ρ , $m_V^2 - m_{PS}^2$, J and f_K/f_π , are insensitive to whether c has its tree-level or tadpole-improved value (Figures 3 and 5).

3. Provided tadpole-improved perturbation theory for Z_A is reliable, there is an indication that f_π/m_ρ may scale for $\beta \geq 6.0$ (Figure 4).

REFERENCES

1. G.P. Lepage and P.B. Mackenzie, Phys. Rev. D48 (1993) 2250.
2. K. Jansen et al. (Alpha), Phys. Lett. B372 (1996) 275.
3. G. Heatlie et al., Nucl. Phys. B352 (1991) 266.
4. F. Butler et al. (GF11), Nucl. Phys. B421 (1994) 217; ibid. B430 (1994) 179.
5. P. Lacock et al. (UKQCD), Phys. Rev. D51 (1995) 6403.
6. C. Michael and H. Shanahan (UKQCD), Nucl. Phys. B (Proc. Suppl.) 47 (1996) 337.
7. R. Gupta and T. Bhattacharya, Los Alamos preprint LAUR-96-1840, hep-lat/9605039.
8. P. Lacock and C. Michael, Phys. Rev. D52 (1995) 5213.

9. P.A. Rowland (UKQCD), these proceedings.




Figure 5. f_K/f_π versus lattice spacing.